

CARE INSTRUCTIONS

TOOTH EXTRACTION

Stop bleeding

- It is important for a blood clot to form to stop the bleeding and begin the healing process.
- Bite on a gauze pad for 30 minutes after the appointment. If the bleeding or oozing persists, place another gauze pad and bite firmly for another 30 minutes. You may have to do this several times.
- After the blood clot forms, it is important not to disturb or dislodge the clot as it aids healing; therefore - Do not rinse vigorously, suck on straws, smoke, drink alcohol or brush teeth next to the extraction site for 72 hours. These activities can dislodge or dissolve the clot and slow down the healing process.
- Limit vigorous exercise for the next 24 hours as this will increase blood pressure and may cause more bleeding from the extraction site.

Pain and Swelling

- An ice pack or an unopened bag of frozen peas or corn applied to the area, will keep swelling to a minimum. The swelling usually subsides after 48 hours.
- Use pain medication as directed.
- If antibiotics are prescribed, continue to take them for the indicated length of time, even if signs and symptoms of infection are gone.
- Drink lots of fluid and eat nutritious soft food on the day of the extraction. You can eat normally as soon as you are comfortable.
- It is important to resume your normal dental routine after 24 hours. This should include brushing and flossing your teeth at least once a day. This will speed up healing and help keep your mouth fresh and clean. You can also rinse with warm salty water: ½ cup of water with ½ teaspoon salt, rinse for 30sec.
- After a few days you will feel fine and can resume your normal activities.

If you have heavy bleeding, severe pain, continued swelling for 2-3 days, or a reaction to the medication, please call our office immediately at (301) 493 – 9500